

1. (12%) Determine the following limit.

(a) (3%)  $\frac{x^2+2x-3}{x^2-1}$

(b) (3%)  $\frac{x^2-4}{|x-2|}$

(c) (3%)  $\frac{3}{1+\frac{2}{x}}$

(d) (3%)  $\frac{\cos(\pi x)}{x+1}$

**Ans:**

(a)  $\frac{x^2+2x-3}{x^2-1} = \frac{(x+3)(x-1)}{(x+1)(x-1)} = \frac{(x+3)}{(x+1)} = 2$

(b)  $\lim_{x \rightarrow 2^+} \frac{x^2-4}{x-2} = \lim_{x \rightarrow 2^+} x + 2 = 4$  and  $\lim_{x \rightarrow 2^-} \frac{x^2-4}{-(x-2)} = \lim_{x \rightarrow 2^-} -(x+2) = -4$

Therefore, the limit does not exist.

(c)  $\frac{3}{1+\frac{2}{x}} = \frac{3}{\frac{x+2}{x}} = \frac{3x}{x+2} = 0$

(d) Since  $\frac{-1}{x+1} \leq \frac{\cos(\pi x)}{x+1} \leq \frac{1}{x+1}$  and  $\frac{1}{x+1} = 0 = \frac{-1}{x+1}$ . By the squeeze theorem,  $\frac{\cos(\pi x)}{x+1} = 0$ .

2. (6%)

If  $f(x)$  and  $g(x)$  are both continuous function with  $[3f(x) + g(x)] = 4$  and  $[f(x) - 2g(x)] = 6$ . Find

(a) (2%)  $f(x)$  (b) (2%)  $g(2)$  (c) (2%)  $f(x)g(x)$

**Ans:**

Since  $f(x)$  and  $g(x)$  are continuous at  $x = 2$ , we have:

$f(x) = f(2)$  and  $g(x) = g(2)$

From the given limits, let  $L = f(2)$  and  $M = g(2)$

$3L + M = 4, L - 2M = 6 \rightarrow L = 2, M = -2$

(a)  $f(x) = L = 2$

(b)  $g(2) = M = -2$

(c)  $f(x)g(x) = -4$

3. (10%)

Let  $f(x) = \begin{cases} \sin(3x) & \text{for } x \leq 0 \\ mx & \text{for } x > 0 \end{cases}$

(a) (5%) Find all values of  $m$  that make  $f$  continuous at 0

(b) (5%) Find all the values of  $m$  that make  $f$  differentiable at 0

**Ans:**

(a)

$f(x) = 0 = f(0) \rightarrow m$  can be any real number.

(b) Considering the alternative form of derivative:

$$\lim_{x \rightarrow 0^-} \frac{f(x) - f(0)}{x - 0} = \lim_{x \rightarrow 0^-} \frac{\sin(3x)}{x} = \lim_{x \rightarrow 0^-} \frac{3\sin(3x)}{3x} = 3 \lim_{t \rightarrow 0^-} \frac{\sin(t)}{t} = 3$$

$$\lim_{x \rightarrow 0^+} \frac{f(x) - f(0)}{x - 0} = \lim_{x \rightarrow 0^+} \frac{mx}{x} = m$$

Since it is differentiable, we have  $m = 3$

4. (5%) If  $f(x) = x^2 + 2x - 3$ , use the definition of the derivative of a function to compute  $f'(x)$

**Ans:**

$$\begin{aligned} f'(x) &= \lim_{\Delta x \rightarrow 0} \frac{(x + \Delta x) - f(x)}{\Delta x} = \lim_{\Delta x \rightarrow 0} \frac{(x + \Delta x)^2 + 2(x + \Delta x) - 3 - (x^2 + 2x - 3)}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} \frac{2x\Delta x + (\Delta x)^2 + 2\Delta x}{\Delta x} = \lim_{\Delta x \rightarrow 0} 2x + \Delta x + 2 = 2x + 2 \end{aligned}$$

5. (5%) Verify that  $f(x) = x^3 + 2x + 4$  satisfies the hypotheses of the Mean Value Theorem on  $[-1,1]$ . Then find all numbers  $c$  that satisfy the conclusion of the Mean Value Theorem.

**Ans:**

The function  $f(x) = x^3 + 2x + 4$  is a polynomial function. Polynomial functions are continuous everywhere on  $R$ , including the closed interval  $[-1,1]$ . Again, since  $f(x)$  is a polynomial function, it is differentiable everywhere on  $R$ , including the open interval  $(-1,1)$ . Therefore, the hypotheses of the Mean Value Theorem are satisfied.

$$f(1) = 7, f(-1) = 1$$

In addition,  $f'(x) = 3x^2 + 2$

By MVT, we have  $f'(c) = \frac{f(1)-f(-1)}{1-(-1)} = 3 \rightarrow 3c^2 + 2 = 3 \rightarrow 3c^2 = 1 \rightarrow c = \pm \frac{\sqrt{3}}{3}$

Both of them lies in  $(-1,1)$ . Therefore,  $c = \pm \frac{\sqrt{3}}{3}$ .

6. (12%)

(a) (5%) Find the equation of the tangent line to the graph of  $f(x) = \frac{x+8}{\sqrt{3x+1}}$  at the point  $(0,8)$

(b) (3%) Use chain rule to find the derivative of  $g(x) = \sin(2x^2 + 3\cos(x))$

(c) (4%) Use implicit differentiation to find  $\frac{dy}{dx}$  and  $\frac{d^2y}{dx^2}$  of the expression  $2xy - 1 = 3x + y^2$

**Ans:**

$$(a) f'(x) = \frac{(3x+1)^{\frac{1}{2}}(1) - (x+8)\frac{1}{2}(3x+1)^{-\frac{1}{2}}(3)}{3x+1}$$

$$f'(0) = \frac{1-4(3)}{1} = -11. \text{ The tangent line is } y - 8 = -11(x - 0) \rightarrow y = -11x + 8$$

$$(b) g'(x) = \cos(2x^2 + 3\cos(x)) \times (4x - 3 \sin(x))$$

(c) Differentiate both side with respect to  $x$ , we have

$$2y + 2x \frac{dy}{dx} = 3 + 2y \frac{dy}{dx} \rightarrow \frac{dy}{dx} = \frac{3 - 2y}{2x - 2y}$$

$$\begin{aligned} \frac{d^2y}{dx^2} &= \frac{-2 \frac{dy}{dx} (2x - 2y) - 2(3 - 2y)(1 - \frac{dy}{dx})}{4(x - y)^2} = \frac{(-4x + 6) \frac{dy}{dx} + (4y - 6)}{4(x - y)^2} \\ &= \frac{(-4x + 6) \left( \frac{3 - 2y}{2x - 2y} \right) + (4y - 6)}{4(x - y)^2} = \frac{-12x + 8xy + 9 - 4y^2}{4(x - y)^3} \end{aligned}$$

7. (20%) Let  $f(x) = \frac{x^4 + x^2 + 4x}{x}$

(a) (4%) Find the critical points and possible points of inflection for  $f(x)$

(b) (3%) Find the open intervals on which  $f(x)$  is increasing or decreasing

(c) (3%) Find the open intervals of concavity for  $f(x)$

(d) (4%) Find all asymptotes of  $f(x)$

(e) (6%) Sketch the graph of  $f(x)$ , labeling intercepts, relative extrema, points of inflection, and asymptotes.

**Ans:** Note that the original function is undefined at  $x = 0$ , therefore we should include it in the following table.

$$f(x) = x^3 + x + 4, \quad x \neq 0, \quad f'(x) = 3x^2 + 1 > 0$$

$$f''(x) = 6x$$

	$(-\infty, 0)$	$(0, \infty)$
測試值	-1	1
$f'$ 的正負號	+	+
$f''$ 的正負號	-	+
結論	遞增/向下凹	遞減/上下凹

(a) Note that  $x$  is not define at  $x = 0$ , we should not include it in the critical numbers or possible points of inflection

There is no critical numbers ( $f' = 0$ )


There is no possible points of inflection ( $f'' = 0$ )

(b) Increasing  $(-\infty, 0), (0, \infty)$ .

(c) Upward:  $(0, \infty)$ . Downward  $(-\infty, 0)$

(d) Since  $f(x) = \pm\infty \rightarrow$  No horizontal asymptote. No vertical asymptote since  $f(x)$  is undefined at  $x = 0$

(e) Graph



There is no relative extrema or point of inflection. No  $y$  intercept

Using Newton's method or bisection method, we have  $x$  intercept roughly equals -1.38. (Other approximation methods are also acceptable, like trial and error)

8. (8%)

(a) (4%) Find the point on the graph  $y = \sqrt{x - 8}$  that is closest to the point  $(12, 0)$ .

(b) (4%) Use Newton's method with the initial approximation  $x_1 = -1$  to find  $x_3$ , the third approximation to the solution of the equation  $2x^3 - 3x^2 + 2 = 0$

**Ans:**

(a) The distance between  $(12, 0)$  and a point  $(x, y)$  on the graph of  $\sqrt{x - 8}$  is

$$d = \sqrt{(x - 12)^2 + y^2} = \sqrt{(x - 12)^2 + x - 8}$$

Minimize  $d^2 = f(x) = (x - 12)^2 + x - 8$ . Note that  $f'(x) = 2(x - 12) + 1 =$

$2x - 23$ . The only critical point is  $x = \frac{23}{2}$ . When  $x = \frac{23}{2}$ ,  $y = \sqrt{\frac{7}{2}}$ ,  $d = \frac{\sqrt{15}}{2}$ . Therefore,

$x = \frac{23}{2}$  is relative minimum (Testing the critical number using First Derivative Test).

Therefore, the closest point is  $(\frac{23}{2}, \sqrt{\frac{7}{2}})$ .

(b) Newton's Method iteratively improves an estimate  $x_n$  of a root of a function  $f(x)$  using the formula:

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)}$$

$$f'(x) = 6x^2 - 6x$$

$n$	$x_n$	$f(x_n)$	$f'(x_n)$	$\frac{f(x_n)}{f'(x_n)}$	$x_n - \frac{f(x_n)}{f'(x_n)}$
1	-1	-3	12	-0.25	-0.75
2	-0.75	-0.53125	7.875	0.06746031746	-0.68254
3	-0.68254				

9. (8%) Use differentials to approximate  $\sqrt{1 + \sin(0.01)}$

**Ans:** Let  $f(x) = \sqrt{1 + \sin(x)}$ ,  $f'(x) = \frac{\cos(x)}{2\sqrt{1 + \sin(x)}}$

$$f(x + \Delta x) \approx f(x) + f'(x)dx = \sqrt{1 + \sin(x)} + \frac{\cos(x)}{2\sqrt{1 + \sin(x)}} dx$$

Choosing  $x = 0$  and  $dx = 0.01$ .

$$f(x + \Delta x) = \sqrt{1 + \sin(0.01)} \approx \sqrt{1 + \sin(0)} + \frac{\cos(0)}{2\sqrt{1 + \sin(0)}} 0.01$$

$$= 1 + \frac{0.01}{2} = 1.005$$

10. (14%) Solve the following problems

(a) (7%) Evaluate  $\int \frac{3}{\sqrt[3]{x}} + x^2 + 2 dx$

(b) (7%) Find  $\frac{1}{n} \left[ \sqrt{\frac{n^2-1^2}{n^2}} + \sqrt{\frac{n^2-2^2}{n^2}} + \sqrt{\frac{n^2-3^2}{n^2}} + \dots + \sqrt{\frac{n^2-n^2}{n^2}} \right]$

**Ans:**

$$(a) \int \frac{3}{\sqrt[3]{x}} + x^2 + 2 dx = \int 3x^{-\frac{1}{3}} + x^2 + 2 dx = \frac{9}{2}x^{\frac{2}{3}} + \frac{1}{3}x^3 + 2x + C$$

$$(b) \left[ \frac{\sqrt{n^2-1^2}}{n^2} + \frac{\sqrt{n^2-2^2}}{n^2} + \frac{\sqrt{n^2-3^2}}{n^2} + \dots + \frac{\sqrt{n^2-n^2}}{n^2} \right] = \sum_{k=1}^n \frac{\sqrt{n^2-k^2}}{n^2} = \sum_{k=1}^n \frac{1}{n} \sqrt{1 - \left(\frac{k}{n}\right)^2} = \int_0^1 \sqrt{1-x^2} dx = \frac{\pi}{4}$$