

Chapter 11 Vectors and the Geometry of Space

Szu-Chi Chung

Department of Applied Mathematics, National Sun Yat-sen University

March 26, 2024

Table of Contents

1 Surfaces in space

2 Cylindrical and spherical coordinates

Table of Contents

1 Surfaces in space

2 Cylindrical and spherical coordinates

Cylindrical surfaces

- You have already known two special types of surfaces.
 - ① Spheres: $(x - x_0)^2 + (y - y_0)^2 + (z - z_0)^2 = r^2$
 - ② Planes: $ax + by + cz + d = 0$
- A third type of surface in space is called a cylindrical surface, or simply a cylinder.
- To define a cylinder, consider the familiar right circular cylinder shown in Figure 1.




Figure 1: Right circular cylinder: $x^2 + y^2 = a^2$. Rulings are parallel to the z -axis.

- You can imagine that this cylinder is generated by a vertical line moving around the circle $x^2 + y^2 = a^2$ in the xy -plane.
- This circle is called a generating curve for the cylinder, as indicated in the following definition.




Figure 2: Right cylinder. Rulings are perpendicular to the plane containing C .

Definition 11.1 (Cylinder)

Let C be a curve in a plane and let L be a line not in a parallel plane. The set of all lines parallel to L and intersecting C is called a cylinder. C is called the generating curve (or **directrix**) of the cylinder, and the parallel lines are called rulings.

- For the right circular cylinder shown in Figure 1, the equation of the generating curve is

$$x^2 + y^2 = a^2. \quad \text{Equation of generating curve in } xy\text{-plane}$$

- To find an equation of the cylinder, note that you can generate any one of the rulings by fixing the values of x and y and then allowing z to take on all real values.
- In this sense, the value of z is arbitrary and is, therefore, not included in the equation. In other words, the equation of this cylinder is simply the equation of its generating curve.

$$x^2 + y^2 = a^2 \quad \text{Equation of cylinder in space}$$


Definition 11.2 (Equation of cylinders)

The equation of a cylinder whose ruling are parallel to one of the coordinate axes contain only the variables corresponding to the other two axes.


Example 1 (Sketching a cylinder)

Sketch the surface represented by each equation.

a. $z = y^2$ **b.** $z = \sin x, 0 \leq x \leq 2\pi$.



(a) Rulings are parallel to the x -axis.



(b) Rulings are parallel to the y -axis.

Quadric surfaces

- The fourth basic type of surface in space is a quadric surface.
Quadric surfaces are the three-dimensional analogs of conic sections.







Definition 11.3 (Quadric surface)






The equation of a quadric surface in space is a second-degree equation in three variables. The general form of the equation is

$$Ax^2 + By^2 + Cz^2 + Dxy + Exz + Fyz + Gx + Hy + Iz + J = 0.$$

There are six basic types of quadric surfaces: ellipsoid, hyperboloid of one sheet, hyperboloid of two sheets, elliptic cone, elliptic paraboloid, and hyperbolic paraboloid.

- The intersection of a surface with a plane is called the trace of the surface in the plane.
- To visualize a surface in space, it is helpful to determine its traces in some well-chosen planes. The traces of quadric surfaces are conics.
- These traces, together with the standard form of the equation of each quadric surface, are shown in the following tables.

	<p style="text-align: center;">Ellipsoid</p> $\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} = 1$ <table border="0" style="width: 100%;"> <tr> <td style="border-bottom: 1px solid black; padding-bottom: 2px;"><i>Trace</i></td> <td style="border-bottom: 1px solid black; padding-bottom: 2px;"><i>Plane</i></td> </tr> <tr> <td>Ellipse</td> <td>Parallel to xy-plane</td> </tr> <tr> <td>Ellipse</td> <td>Parallel to xz-plane</td> </tr> <tr> <td>Ellipse</td> <td>Parallel to yz-plane</td> </tr> </table> <p>The surface is a sphere if $a = b = c \neq 0$.</p>	<i>Trace</i>	<i>Plane</i>	Ellipse	Parallel to xy -plane	Ellipse	Parallel to xz -plane	Ellipse	Parallel to yz -plane	
<i>Trace</i>	<i>Plane</i>									
Ellipse	Parallel to xy -plane									
Ellipse	Parallel to xz -plane									
Ellipse	Parallel to yz -plane									
	<p style="text-align: center;">Hyperboloid of One Sheet</p> $\frac{x^2}{a^2} + \frac{y^2}{b^2} - \frac{z^2}{c^2} = 1$ <table border="0" style="width: 100%;"> <tr> <td style="border-bottom: 1px solid black; padding-bottom: 2px;"><i>Trace</i></td> <td style="border-bottom: 1px solid black; padding-bottom: 2px;"><i>Plane</i></td> </tr> <tr> <td>Ellipse</td> <td>Parallel to xy-plane</td> </tr> <tr> <td>Hyperbola</td> <td>Parallel to xz-plane</td> </tr> <tr> <td>Hyperbola</td> <td>Parallel to yz-plane</td> </tr> </table> <p>The axis of the hyperboloid corresponds to the variable whose coefficient is negative.</p>	<i>Trace</i>	<i>Plane</i>	Ellipse	Parallel to xy -plane	Hyperbola	Parallel to xz -plane	Hyperbola	Parallel to yz -plane	
<i>Trace</i>	<i>Plane</i>									
Ellipse	Parallel to xy -plane									
Hyperbola	Parallel to xz -plane									
Hyperbola	Parallel to yz -plane									
	<p style="text-align: center;">Hyperboloid of Two Sheets</p> $\frac{z^2}{c^2} - \frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$ <table border="0" style="width: 100%;"> <tr> <td style="border-bottom: 1px solid black; padding-bottom: 2px;"><i>Trace</i></td> <td style="border-bottom: 1px solid black; padding-bottom: 2px;"><i>Plane</i></td> </tr> <tr> <td>Ellipse</td> <td>Parallel to xy-plane</td> </tr> <tr> <td>Hyperbola</td> <td>Parallel to xz-plane</td> </tr> <tr> <td>Hyperbola</td> <td>Parallel to yz-plane</td> </tr> </table> <p>The axis of the hyperboloid corresponds to the variable whose coefficient is positive. There is no trace in the coordinate plane perpendicular to this axis.</p>	<i>Trace</i>	<i>Plane</i>	Ellipse	Parallel to xy -plane	Hyperbola	Parallel to xz -plane	Hyperbola	Parallel to yz -plane	
<i>Trace</i>	<i>Plane</i>									
Ellipse	Parallel to xy -plane									
Hyperbola	Parallel to xz -plane									
Hyperbola	Parallel to yz -plane									

	<p style="text-align: center;">Elliptic Cone</p> $\frac{x^2}{a^2} + \frac{y^2}{b^2} - \frac{z^2}{c^2} = 0$ <table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="text-align: left;">Trace</th> <th style="text-align: left;">Plane</th> </tr> </thead> <tbody> <tr> <td>Ellipse</td> <td>Parallel to xy-plane</td> </tr> <tr> <td>Hyperbola</td> <td>Parallel to xz-plane</td> </tr> <tr> <td>Hyperbola</td> <td>Parallel to yz-plane</td> </tr> </tbody> </table> <p>The axis of the cone corresponds to the variable whose coefficient is negative. The traces in the coordinate planes parallel to this axis are intersecting lines.</p>	Trace	Plane	Ellipse	Parallel to xy -plane	Hyperbola	Parallel to xz -plane	Hyperbola	Parallel to yz -plane	
Trace	Plane									
Ellipse	Parallel to xy -plane									
Hyperbola	Parallel to xz -plane									
Hyperbola	Parallel to yz -plane									
	<p style="text-align: center;">Elliptic Paraboloid</p> $z = \frac{x^2}{a^2} + \frac{y^2}{b^2}$ <table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="text-align: left;">Trace</th> <th style="text-align: left;">Plane</th> </tr> </thead> <tbody> <tr> <td>Ellipse</td> <td>Parallel to xy-plane</td> </tr> <tr> <td>Parabola</td> <td>Parallel to xz-plane</td> </tr> <tr> <td>Parabola</td> <td>Parallel to yz-plane</td> </tr> </tbody> </table> <p>The axis of the paraboloid corresponds to the variable raised to the first power.</p>	Trace	Plane	Ellipse	Parallel to xy -plane	Parabola	Parallel to xz -plane	Parabola	Parallel to yz -plane	
Trace	Plane									
Ellipse	Parallel to xy -plane									
Parabola	Parallel to xz -plane									
Parabola	Parallel to yz -plane									
	<p style="text-align: center;">Hyperbolic Paraboloid</p> $z = \frac{y^2}{b^2} - \frac{x^2}{a^2}$ <table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="text-align: left;">Trace</th> <th style="text-align: left;">Plane</th> </tr> </thead> <tbody> <tr> <td>Hyperbola</td> <td>Parallel to xy-plane</td> </tr> <tr> <td>Parabola</td> <td>Parallel to xz-plane</td> </tr> <tr> <td>Parabola</td> <td>Parallel to yz-plane</td> </tr> </tbody> </table> <p>The axis of the paraboloid corresponds to the variable raised to the first power.</p>	Trace	Plane	Hyperbola	Parallel to xy -plane	Parabola	Parallel to xz -plane	Parabola	Parallel to yz -plane	
Trace	Plane									
Hyperbola	Parallel to xy -plane									
Parabola	Parallel to xz -plane									
Parabola	Parallel to yz -plane									

Example 2 (Sketching a quadric surface)

Classify and sketch the surface given by

$$4x^2 - 3y^2 + 12z^2 + 12 = 0.$$




Figure 4: Hyperboloid of two sheets: $\frac{y^2}{4} - \frac{x^2}{3} - z^2 = 1$.

Example 3 (Sketching a quadric surface)

Classify and sketch the surface given by $x - y^2 - 4z^2 = 0$.





Figure 5: Elliptic paraboloid.

Example 4 (A quadric surface not centered at the origin)

Classify and sketch the surface given by
 $x^2 + 2y^2 + z^2 - 4x + 4y - 2z + 3 = 0$.

$$\frac{(x-2)^2}{4} + \frac{(y+1)^2}{2} + \frac{(z-1)^2}{4} = 1$$




Surfaces of revolution

- The fifth special type of surface you will study is called a surface of revolution. We now look at how to find its equation.
- Consider the graph of the radius function

$$y = r(z) \quad \text{Generating curve}$$

in the yz -plane.

- If this graph is revolved around the z -axis, it forms a surface of revolution.



- The trace of the surface in the plane $z = z_0$ is a circle whose radius is $r(z_0)$ and whose equation is

$$x^2 + y^2 = [r(z_0)]^2. \quad \text{Circular trace in plane: } z = z_0$$

- Replacing z_0 with z produces an equation that is valid for all values of z .
- You can obtain equations for surfaces of revolution for the other two axes, and the results are summarized as follows.

Definition 11.4 (Surface of revolution)

If the graph of a radius function r is revolved about one of the coordinate axes, the equation of the resulting surface of revolution has one of the following forms.

- 1 Revolved about the x -axis: $y^2 + z^2 = [r(x)]^2$
- 2 Revolved about the y -axis: $x^2 + z^2 = [r(y)]^2$
- 3 Revolved about the z -axis: $x^2 + y^2 = [r(z)]^2$

Example 5 (Finding an equation for a surface of revolution)

Find an equation for the surface of revolution formed by revolving (a) the graph of $y = 1/z$ about the z -axis and (b) the graph of $9x^2 = y^3$ about the y -axis.




Figure 6: Surface of revolution: $x^2 + z^2 = \frac{1}{9}y^3$ with generating curve $9x^2 = y^3$ about the y -axis.

Example 6 (Finding a generating curve for a surface of revolution)

Find a generating curve and the axis of revolution for the surface given by

$$x^2 + 3y^2 + z^2 = 9.$$




Figure 7: Finding a generating curve for a surface of revolution: not unique.

Table of Contents

1 Surfaces in space

2 Cylindrical and spherical coordinates

Cylindrical coordinates

- The cylindrical coordinate system, is an extension of the polar coordinates in the plane to three-dimensional space.

Definition 11.5 (The cylindrical coordinate system)

In a cylindrical coordinate system, a point P in space is represented by an ordered triple (r, θ, z) .

- 1 (r, θ) is a polar representation of the projection of P in the xy -plane.
- 2 z is the directed distance from (r, θ) to P .

- To convert from rectangular to cylindrical coordinates (or vice versa), use the following conversion guidelines for polar coordinates:

$$x = r \cos \theta, \quad y = r \sin \theta, \quad z = z$$

$$r^2 = x^2 + y^2, \quad \tan \theta = \frac{y}{x}, \quad z = z$$




Figure 8: The relationship between cylindrical and rectangular coordinates.

- The point $(0, 0, 0)$ is called the pole.

- Moreover, because the representation of a point in the polar coordinate system is not unique, it follows that the representation in the cylindrical coordinate system is also not unique!

Example 1 (Converting from cylindrical to rectangular coordinates)

Convert the point $(r, \theta, z) = (4, \frac{5\pi}{6}, 3)$ to rectangular coordinates.




Figure 9: Converting $(r, \theta, z) = \left(4, \frac{5\pi}{6}, 3\right)$ to $(x, y, z) = (-2\sqrt{3}, 2, 3)$.

Example 2 (Converting from rectangular to cylindrical coordinate)

Convert the point $(x, y, z) = (1, \sqrt{3}, 2)$ to cylindrical coordinates.




Figure 10: Converting from rectangular to cylindrical coordinates.

- Cylindrical coordinates are especially convenient for representing cylindrical surfaces and surfaces of revolution with the z-axis as the axis of symmetry:




Figure 11: Different cylindrical equations.

- Vertical planes containing the z -axis and horizontal planes also have simple cylindrical coordinate equations:




Figure 12: Vertical plane: $\theta = c$ and horizontal plane: $z = c$.

Example 3 (Rectangular-to-cylindrical conversion)

Find an equation in cylindrical coordinates for the surface represented by each rectangular equation.

a. $x^2 + y^2 = 4z^2$ **b.** $y^2 = x$






Figure 13: Rectangular-to-cylindrical conversion.

Example 4 (Cylindrical-to-rectangular conversion)

Find an equation in rectangular coordinates for the surface represented by the cylindrical equation

$$r^2 \cos 2\theta + z^2 + 1 = 0.$$

Cylindrical:
 $r^2 \cos 2\theta + z^2 + 1 = 0$



Rectangular:
 $y^2 - x^2 - z^2 = 1$

Figure 14: Cylindrical-to-rectangular conversion.

Spherical coordinates

- In the spherical coordinate system, each point is represented by an ordered triple: the first coordinate is a distance, and the second and third coordinates are angles.
- This system is similar to the latitude-longitude system used to identify points on the surface of Earth.
- For example, the point on the surface of Earth whose latitude is 40° North (of the equator) and whose longitude is 80° West (of the prime meridian) is shown in Figure 15. Assuming that the Earth is spherical and has a radius of 6371 kilometers, you would label this point as

$$(4000, -80^\circ, 50^\circ).$$

Radius 80° clockwise from
prime meridian 50° down from
North Pole




Figure 15: Spherical coordinate of 80° W 40° N is $(4000, -80^\circ, 50^\circ)$.

Definition 11.6 (The spherical coordinate system)

In a spherical coordinate system, a point P in space is represented by an ordered triple (ρ, θ, ϕ) .

1. ρ is the distance between P and the origin, $\rho \geq 0$.
2. θ is the same angle used in cylindrical coordinates for $r \geq 0$.
3. ϕ is the angle between the positive z -axis and the line segment \overrightarrow{OP} , $0 \leq \phi \leq \pi$.

Note that the first and third coordinates, ρ and ϕ , are nonnegative. ρ is the lowercase Greek letter rho, and ϕ is the lowercase Greek letter phi.

- The relationship between rectangular and spherical coordinates is illustrated in Figure 16.




Figure 16: The relationship between rectangular coordinate (x, y, z) and spherical coordinates (ρ, θ, ϕ) where $r = \rho \sin \phi = \sqrt{x^2 + y^2}$.

- To convert from one system to the other, use the following.
- Spherical to rectangular:

$$x = \rho \sin \phi \cos \theta, \quad y = \rho \sin \phi \sin \theta, \quad z = \rho \cos \phi.$$

- Rectangular to spherical:

$$\rho^2 = x^2 + y^2 + z^2, \quad \tan \theta = \frac{y}{x}, \quad \phi = \arccos \left(\frac{z}{\sqrt{x^2 + y^2 + z^2}} \right).$$


- To change coordinates between the cylindrical and spherical systems, use the following.
- Spherical to cylindrical ($r \geq 0$):

$$r^2 = \rho^2 \sin^2 \phi, \quad \theta = \theta, \quad z = \rho \cos \phi.$$


- Cylindrical to spherical ($r \geq 0$):

$$\rho = \sqrt{r^2 + z^2}, \quad \theta = \theta, \quad \phi = \arccos \left(\frac{z}{\sqrt{r^2 + z^2}} \right).$$


- The spherical coordinate system is useful primarily for surfaces in space that have a point or center of symmetry.
- For example, Figure 17 shows three surfaces with simple spherical equations.



Sphere:
 $\rho = c$



Vertical half-plane:
 $\theta = c$



Half-cone: $\left(0 < c < \frac{\pi}{2}\right)$
 $\phi = c$

Figure 17: Three surfaces with simple spherical equations.

Example 5 (Rectangular-to-spherical conversion)

Find an equation in spherical coordinates for the surface represented by each rectangular equation.

a. Cone: $x^2 + y^2 = z^2$ **b.** Sphere: $x^2 + y^2 + z^2 - 4z = 0$




Figure 18: $x^2 + y^2 + z^2 - 4z = 0$ in rectangular coordinate is equivalent to $\rho = 4 \cos \phi$ in spherical coordinate.